Immunoglobulin Referral Form

New to therapy Therapy Continuat	ion			
Deliver to: Patient's home Prescr		Date Initiated:	Date Needed:	
PATIENT INFORMATION		PRESCRIBER INFORM	ATION	
Full name				
	Male 🗌 Female	State License		
Street address		NPI		
City	State Zip	Address		
	Secondary phone			
Patient's guardian	HIPPA Consent [] Yes [] No	Fax		
		CLINICAL INFORMATIO	N	
		-	t: Date recorded:	
			gies	
	Group	Previous Immunoglobulin Ther	-	
Does patient have secondary insurance?			apies (ii applicable).	
	Please provide copy of primary an	d secondary insurance with this fo		
	riease provide copy of primary an	a secondary insurance with this ic	лш 	
PRESCRIBING INFORMATION				
No Brand Preference	Brand Products		Other	
☐ Immune Globulin Solution ☐ 5% ☐ 10% ☐ 20%	☐ Bivigam 10% ☐ Cuvitru 20%	Gamunex-C 10%	(Specify Product)	
(No Brand Preference)	☐ Flebogamma			
	5% 10%	☐ Hyqvia 10%		
	Gammagard Liquid 10%	☐ Octagam		
	Gammagard S/D low IgA	5% <u></u> 10%		
	☐ 5% ☐ 10%	Panzyga 10%		
	Gammaked 10%	Privigen 10%		
Route: IV SQ	Qty:Directions:			
IV access:		Pre/Post Hydration:		
Peripheral Port PICC		mL of 0.9% NaCl D5W before after concurrently at a rate of		
Flush Protocol:		Other:	mL/hour.	
Use 5mL to 10mL of 0.9% NaCl before and after each infusion. Sterile syringes required for PICC/PORT.		Anaphylxis Kit:		
Maintain PICC with 3 to 5mL of 10unit/mL of heparin and maintain implanted port with 3 to 5mL		IVIG: Provide anaphylaxis kit per protocol (epinephrine 1 mg/mL ampule, diphenhydramine		
of 100unit/mL of heparin.		50 mg/mL vial, diphenhydramine 12.5mg or 25mg tablets or capsules, 1000cc 0.9% NaCl, all infusion supplies)		
Pre-medication:		SQIG: Epinephrine Pen 2-pack (0.3 mg for ≥30 kg; 0.15 mg for <30 kg) Sig: Inject IM in		
Acetaminophen 325mg tablets Sig: Take two 325mg tablets (650mg) by mouth 30-60 minutes prior to infusion.		event of anaphylaxis Qty: 1 pack Refills: PRN		
Qty: 2 per dose		Nursing Care:		
Diphenhydramine 25mg capsules		Infused in office or infusion center Home Nursing needed Nursing already coordinated:		
Sig: Take one to two 25mg capsules (25-50mg) by mouth 30-60 minutes prior to infusion. Qtv: 2 per dose		Agency		
		RN to provide home nursing services for administration of IVIG or patient teach of SQIG,		
Medications to be used as needed:		and as needed for IV site care and complications related to therapy.		
Sig: Apply small amount to injection site 60 min prior to infusion		Supplies: Pharmacy will provide home infusion.	all supplies, fluids and ancillary equipment necessary for	
Qty: 1 tube	0	nome musion.		
	Qty:			
	supply or please specify if other:			
Substitution Permissible. In order for a	brand name product to be dispensed, the prescriber m	ust handwrite "BRAND NECESSARY		
Prescriber's Signature (Dispense as Written)				
Prescriber's Signature (Substitution Pe	ermissible)		Date:	

For ARNP, NP, and PA, collaborative physician agreement is with: _

CONFIDENTIAL HEALTH INFORMATION: Healthcare information is personal information related to a person's healthcare. It is being faxed to you after appropriate authorization or under circumstances that don't require authorization. You are obligated to maintain it in a safe, secure, and confidential manner. Re-disclosure of this information is prohibited unless permitted by law or appropriate customer/patient authorization is obtained. Unauthorized re-disclosure or failure to maintain confidentiality could subject you to penalties described in federal and state laws. Drug names are the property of their respective owners.